

Analytical expressions of the dispersive contributions of the nuclear optical model

J.M. Quesada,* A. Molina, and M. Lozano
*Departamento de Física Atómica, Molecular y Nuclear,
 Universidad de Sevilla, Ap. 1065, E-41080 Sevilla, Spain*

R. Capote†
Departamento de Física Aplicada, Universidad de Huelva, E-21071 Huelva, Spain

J. Raynal
*4 rue du Bief, 91380 Chilly-Mazarin, France
 (Dated: February 8, 2008)*

Analytical solutions of dispersion relations in the nuclear optical model have been found for both imaginary volume and surface potentials. A standard Brown-Rho shape has been assumed for the volume imaginary term and a Brown-Rho shape multiplied by a decreasing exponential for the surface contribution. The analytical solutions are valid for any even value of the exponent appearing in these functional forms.

PACS numbers: 11.55.Fv, 24.10.Ht

A significant contribution to the optical model theory during the last two decades can be considered the work of Mahaux and co-workers on dispersive optical model analysis [1, 2, 3, 4, 5]. The unified description of nuclear mean field in dispersive optical model is accomplished by using a dispersion relation, which links the real and absorptive terms of the optical model potential (OMP). The additional constraint imposed by dispersion relation helps to reduce the ambiguities in deriving phenomenological OMP parameters from the experimental data. In a dispersion relation treatment, the real central potential strength consists of a term which varies slowly with energy E , the so called Hartree-Fock (HF) term, $V_{HF}(r, E)$, plus a correction term, $\Delta V(r, E)$, which is calculated using a dispersion relation. The depth of the dispersive term of the potential $\Delta V(r, E)$ can be written as

$$\Delta V(r, E) = \frac{\mathcal{P}}{\pi} \int_{-\infty}^{\infty} \frac{W(r, E')}{E' - E} dE' \quad (1)$$

Being $W(r, E)$ the imaginary part of the OMP. Assuming that $W(r, E = E_F) = 0$ and $\Delta V(r, E = E_F) = 0$, where E_F is the Fermi energy, then equation (1) can be written in the subtracted form:

$$\Delta V(r, E) = \frac{\mathcal{P}}{\pi} \int_{-\infty}^{\infty} W(r, E') \left(\frac{1}{E' - E} - \frac{1}{E' - E_F} \right) dE' \quad (2)$$

With the assumption that $W(r, E)$ be symmetric respect to the Fermi energy, the equation (2) can be expressed in a form which is stable under numerical treatment [6],

namely:

$$\Delta V(r, E) = \frac{2}{\pi} (E - E_F) \int_{E_F}^{\infty} \frac{W(r, E') - W(r, E)}{(E' - E_F)^2 - (E - E_F)^2} dE' \quad (3)$$

The dispersive term $\Delta V(r, E)$ is divided into two terms $\Delta V_v(r, E)$ and $\Delta V_s(r, E)$, which arise through dispersion relations (1) from the volume $W_v(r, E)$ and surface $W_s(r, E)$ imaginary potentials respectively. If the imaginary potential geometry is energy dependent, the radial dependence of the dispersive correction can not be expressed using a Woods-Saxon form factor $f_{WS}(r)$. However, to simplify the problem, the OMP geometry parameters are usually assumed to be energy independent. In this case, $\Delta V(r, E) = \Delta V(E) f_{WS}(r)$ and the energy dependence of the real volume $V_v(E)$ and surface $V_s(E)$ parts of the dispersive OMP are given by:

$$\begin{aligned} V_v(E) &= V_{HF}(E) + \Delta V_v(E) \\ V_s(E) &= \Delta V_s(E) \end{aligned} \quad (4)$$

It is useful to represent the variation of surface $W_s(E)$ and volume $W_v(E)$ absorption potential depths with energy in functional forms suitable for the dispersive optical model analysis. An energy dependence for the imaginary volume term has been suggested in studies of nuclear matter theory [7]:

$$W_v(E) = A_v \frac{(E - E_F)^n}{(E - E_F)^n + (B_v)^n} \quad (5)$$

where A_v and B_v are undetermined constants. Brown and Rho [7], propose $n = 2$, while Mahaux and Sartor [2] have suggested $n = 4$.

An energy dependence for the imaginary-surface term has been suggested by Delaroche *et al* [6] to be:

$$W_s(E) = A_s \frac{(E - E_F)^m}{(E - E_F)^m + (B_s)^m} \exp(-C_s |E - E_F|) \quad (6)$$

*quesada@us.es

†Permanent address: Centro de Estudios Aplicados al Desarrollo Nuclear, Ap. 100, Miramar, La Habana, Cuba

where $m = 2, 4$ and A_s, B_s and C_s are undetermined constants.

According to equations (5) and (6) the imaginary part of the OMP is assumed to be zero at $E = E_F$ and nonzero everywhere else. A more realistic parameterization of $W_v(E)$ and $W_s(E)$ forces these terms to be zero in some region around the Fermi energy. A physically reasonable energy for defining such a region is the average energy of the single-particle states E_p [4].

$$W_s(E) = \begin{cases} 0 & E_F < E < E_p \\ A_s e^{-C_s |E-E_p|} \frac{(E-E_p)^m}{(E-E_p)^m + (B_s)^m} & E \geq E_p \end{cases} \quad (8)$$

The symmetry condition

$$W(2E_F - E) = W(E) \quad (9)$$

is used to define imaginary part of the OMP for energies below the Fermi energy. Equations (7-9) are used to describe imaginary absorptive potential in this contribution.

Nonlocality effects on the imaginary part can be treated according to the work of Mahaux and Sartor [4] and the analytical solutions of their contributions coming from the dispersion relations have been found by VanderKam *et al.* [8].

In a recent work we have presented a general numerical solution of the dispersion integral relations between the real and the imaginary parts of the nuclear optical potential [9]. In this work we develop analytical solutions of dispersion relations, equation (2) for usually employed functional forms of the imaginary potential, $W(E)$, used in dispersive optical model analyses (equations (7-9)). According to [8] we introduce the quantities $E_0 = E_p - E_F$ (the offset energy), $E_+ = E_x + E_0$ and $E_- = E_x - E_0$. U and E_x are the integration variable and the excitation energy ($E - E_F$), respectively. In the next formulae, we will drop the subscripts on the constants A, B, C .

For the surface potential $W_s(E)$, given by equations (8) and (9), we can write the dispersive integral (2) for m even as

$$\begin{aligned} \Delta V_s(E) &= \frac{E_x}{\pi} \mathcal{P} \int_{-\infty}^{\infty} \frac{W_s(E')}{(E' - E)(E' - E_F)} dE' = \\ &= \frac{E_x}{\pi} \mathcal{P} \int_0^{\infty} \frac{U^m \exp(-CU)}{(U^m + B^m)(U - E_-)(U + E_0)} dU + \\ &+ \frac{E_x}{\pi} \mathcal{P} \int_0^{\infty} \frac{U^m \exp(-CU)}{(U^m + B^m)(U + E_+)(U + E_0)} dU \quad (10) \end{aligned}$$

and the same expression without exponential for the volume potential $W_v(E)$, given by equations (7) and (9).

Therefore a new definition for imaginary part of the OMP can be written as:

$$W_v(E) = \begin{cases} 0 & E_F < E < E_p \\ A_v \frac{(E-E_p)^n}{(E-E_p)^n + (B_v)^n} & E \geq E_p \end{cases} \quad (7)$$

and likewise for surface absorption.

The integrand can be replaced by its expressions in terms of poles and residues:

$$\begin{aligned} &\frac{E_x}{\pi} \frac{U^m}{(U^m + B^m)(U \mp E_{\mp})(U + E_0)} = \\ &= \frac{1}{\pi} \left\{ \sum_{j=1}^m \frac{Res(p_j)}{U - p_j} + \frac{Res(\mp E_{\pm})}{U \mp E_{\mp}} + \frac{Res(E_0)}{U + E_0} \right\} \quad (11) \end{aligned}$$

where the p_j are the m zeroes of $(U^m + B^m)$ and the $Res(p_j)$ their residue, that is :

$$p_j = B \exp\left(i \frac{2j-1}{m} \cdot \pi\right) \quad (12)$$

$$Res(p_j) = \frac{E_x}{m} \frac{p_j}{(p_j \mp E_{\mp})(p_j + E_0)} \quad (13)$$

where $\pm E_{\mp}$ and $-E_0$ are the poles of $U \mp E_{\mp}$ and $U + E_0$, whereas $Res(\pm E_{\mp})$ and $Res(-E_0)$ are their residues respectively

$$Res(\pm E_{\mp}) = \pm \frac{(E_{\mp})^m}{(E_{\mp})^m + B^m} \quad (14)$$

$$Res(-E_0) = \mp \frac{(E_0)^m}{(E_0)^m + B^m} \quad (15)$$

As was pointed out by Raynal [10] the contribution of each complex pole p_j to the surface dispersive integral is :

$$\begin{aligned} &\int_0^{\infty} \frac{Res(p_j) e^{-CU}}{U - p_j} dU = \\ &= Res(p_j) e^{-Cp_j} \int_{-Cp_j}^{\infty} \frac{\exp(-z)}{z} dz \equiv \\ &\equiv Res(p_j) e^{-Cp_j} E_1(-Cp_j) \quad (16) \end{aligned}$$

where $E_1(z)$ is the *Exponential Integral Function* E_1 [11]. For the real poles corresponding to the second term in the

right side of (11), the contribution of each one is given by:

$$\begin{aligned}
& \int_0^\infty \frac{Res(\mp E_\pm) e^{-CU}}{U \mp E_\mp} dU = \\
& = Res(\mp E_\pm) e^{\mp CE_\mp} \mathcal{P} \int_{\mp CE_\mp}^\infty \frac{\exp(-x)}{x} dx \equiv \\
& \equiv -Res(\mp E_\pm) e^{\mp CE_\mp} Ei(\pm CE_\mp) \quad (17)
\end{aligned}$$

where $Ei(x)$ is the *Exponential Integral Function* Ei [11]. The contributions coming from the third term of the right side of eq.(11) in the integral (10) cancel.

For the volume dispersive integral the exponential is missed in the integrand, therefore each pole gives a divergent contribution. Nevertheless their sum is a finite quantity, which can be calculated by taking the proper limit.

We are quoting below exact expressions for the surface and volume dispersive integrals for any even values of m and n in the surface and volume imaginary potentials respectively. This make a difference to ref. [8], where, in surface dispersive integral with $m = 4$, approximations were made limiting the energy range of validity for given analytical expressions.

For the case of the dispersion relation using surface imaginary potential $W_s(E)$ with m even according to equations (8) and (9), the dispersive correction is

$$\begin{aligned}
\Delta V_s(E) = & \frac{A}{\pi} \left\{ \sum_{j=1}^m Z_j e^{-p_j C} E_1(-p_j C) \right. \\
& \left. - Res(-E_+) e^{CE_+} Ei(-CE_+) \right\}
\end{aligned}$$

$$- Res(E_-) e^{-CE_-} Ei(CE_-) \Big\} \quad (18)$$

where Z_j comes from the sum of residues $Res(p_j)$ in the two integrals (10) and is given by

$$Z_j = \frac{E_x}{m} \frac{p_j(2p_j + E_+ - E_-)}{(p_j + E_0)(p_j + E_+)(p_j - E_-)} \quad (19)$$

For the case of the dispersion relation using the volume imaginary potential $W_v(E)$ with n even according to equations (7) and (9), the dispersive correction is

$$\begin{aligned}
\Delta V_v(E) = & -\frac{A}{\pi} \left\{ \sum_{j=1}^n Z_j \ln(-p_j) \right. \\
& + Res(-E_+) \ln E_+ \\
& \left. + Res(E_-) \ln |E_-| \right\} \quad (20)
\end{aligned}$$

In conclusion, we have developed analytical solutions of dispersion relations for the volume and surface terms of the OMP. The formulae are compact and easy to implement in current generation of the optical model parameter search codes. We stress that the solutions are valid for any even exponent n or m .

This work was supported by Junta de Andalucía and the Spanish CICYT under Contracts PB1998-1111, FPA2001-0144-C05-03 and FPA2001-4960-E and by the European Union under Contract FKIW-CT-2000-00107.

-
- [1] C. Mahaux and H. Ngô, Nucl Phys. **A431**, 486 (1984).
 - [2] C. Mahaux and R. Sartor, Nucl Phys. **A468**, 193 (1987).
 - [3] C. H. Johnson, D. J. Horen, and C. Mahaux, Phys. Rev. C **36**, 2252 (1987).
 - [4] C. Mahaux and R. Sartor, Nucl. Phys. **A528**, 253 (1991).
 - [5] C. Mahaux and R. Sartor, *Advances in Nuclear Physics*, vol. 20 ((edited by J.W. Negele and E. Vogt. Plenum, New York), 1991).
 - [6] J. P. Delaroche, Y. Wang, and J. Rapaport, Phys. Rev. C **39**, 391 (1989).
 - [7] G. E. Brown and M. Rho, Nucl Phys. **A372**, 397 (1981).
 - [8] J. M. VanderKam, G. J. Weisel, and W. Tornow, J. Phys. G:Nucl.Part.Phys. **26**, 1787 (2000).
 - [9] R. Capote, A. Molina, and J. M. Quesada, J. Phys. G:Nucl.Part.Phys. **27**, B15 (2001).
 - [10] J. Raynal, in *Proceedings of the Specialists' Meeting on the Nucleon Nucleus Optical Model up to 200 MeV*, Bruyères-le-Châtel, 1996. Available at <http://www.nea.fr/html/science/om200/>. (1997).
 - [11] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions, Applied Mathematics Series, vol. 55* ((Washington: National Bureau of Standards; reprinted by Dover Publications, New York), 1968).